



Published in final edited form as:

J Neurosci. 2009 July 15; 29(28): 8991–8995. doi:10.1523/JNEUROSCI.5717-08.2009.

Neocortical disynaptic inhibition requires somatodendritic integration in interneurons

Court Hull, Hillel Adesnik, and Massimo Scanziani

Neurobiology Section, Division of Biology, University of California at San Diego, La Jolla, California 92093-0634

Abstract

In his theory of functional polarity Ramon y Cajal first identified the soma and dendrites as the principal recipient compartments of a neuron and the axon as its main output structure. Despite notable exceptions in other parts of the nervous system (Schoppa and Urban, 2003; Wassle, 2004; Howard et al., 2005), this route of signal propagation has been shown to underlie the functional properties of most neocortical circuits studied so far. Recent evidence, however, suggests that neocortical excitatory cells may trigger the release of the inhibitory neurotransmitter GABA by directly depolarizing the axon terminals of inhibitory interneurons, thus bypassing their somatodendritic compartments (Ren et al., 2007). By using a combination of optical and electrophysiological approaches, we find that synaptically released glutamate fails to trigger GABA release through a direct action on GABAergic terminals under physiological conditions. Rather, our evidence suggests that glutamate triggers GABA release only following somatodendritic depolarization and action potential generation at GABAergic interneurons. These data indicate that neocortical inhibition is recruited by classical somatodendritic integration rather than direct activation of interneuron axon terminals.

Keywords

Synapse; Inhibition; GABA; Cortex; Somatosensory cortex; Visual

Introduction

The general model of information transfer in the central nervous system involves electrochemical communication between the axon terminal of a presynaptic cell and the somatodendritic compartments of a postsynaptic target cell. In this model, the soma and dendrites of a postsynaptic neuron integrate its excitatory and inhibitory synaptic inputs to produce an output when the membrane depolarization is sufficient to trigger an action potential. There are neuronal circuits, however, which bypass this type of somatodendritic integration. In the spinal cord, for example, glutamate release from sensory afferents is regulated by inhibitory synapses located on the afferent's axon terminal (Rudomin, 1990). Furthermore, in the thalamus, excitatory afferent inputs target the specialized dendrites of inhibitory interneurons, which are capable of directly releasing GABA as a result of local depolarization rather than somatodendritic integration and action potential generation (Sherman, 2004).

Correspondence should be addressed to: Massimo Scanziani Howard Hughes Medical Institute University of California, San Diego Neurobiology Section 0634 Division of Biology 9500 Gilman Drive, La Jolla, CA 92093-0634, U.S.A Phone (Office): (858)-822-3839 Phone (Lab): (858)-822-3840 Fax: (858)-534-2890 e-mail: massimo@biomail.ucsd.edu.

Recent evidence from the visual cortex suggests that axons from layer 2/3 pyramidal cells may directly contact the axon terminals of inhibitory interneurons, thus enabling the pyramidal cells to trigger GABA release onto their neighbors while bypassing somatodendritic integration in interneurons (Ren et al., 2007). This would allow selective and localized output from only part of an interneuron's axonal arborization. Interneurons would hence provide localized inhibition as specified by the anatomical pattern of pyramidal cell synapses onto their axon terminals rather than global inhibition to their cumulative target population. This type of circuit, however, contrasts with the lack of anatomical evidence for glutamatergic axo-axonic contacts in the neocortex (for discussion, see Connors and Cruikshank, 2007). Here, we show that the observation suggesting a direct action of glutamate onto GABAergic terminals can be in part accounted for by an increased glutamate release onto the somatodendritic compartment of GABAergic interneurons using a cesium-based internal solution. Furthermore, we show that release of glutamate by direct activation of channelrhodopsin-expressing axon terminals does not drive GABAergic terminals to release GABA. We therefore find no evidence that pyramidal cells can directly activate the presynaptic terminals of neocortical interneurons to trigger GABA release under normal conditions, and conclude that neocortical inhibition is recruited by classical somatodendritic integration in these circuits.

Methods

Slice preparation

Acute coronal slices of the primary visual and somatosensory cortices (300 μm) were prepared from ICR white mice (p14-27) and Wistar rats (p22-30). Mice were anesthetized with isoflurane, and rats were anesthetized with ketamine/xylazine. For experiments involving recordings from interneurons, G42 and B13 (p14-30) mice were used (Chattopadhyaya et al., 2004; Dumitriu et al., 2007). Slices were incubated on an interface chamber for 30 min at 35 $^{\circ}$ C with an artificial cerebrospinal fluid equilibrated with 95% O₂ and 5% CO₂, containing (in mM): 126 NaCl, 3 KCl, 1.2 NaH₂PO₄, 1.3 MgSO₄, 2.4CaCl₂, 26 NaHCO₃, and 10 glucose (pH 7.3, osmolarity 310). Following incubation, slices were kept in the same chamber at room temperature for 0-6 hr until being transferred to a submerged chamber where electrophysiological recordings were made at 31-33 $^{\circ}$ C.

Recordings

Whole-cell recordings of visually identified pyramidal cells in layer 2/3 (infrared DIC videomicroscopy and water immersion 40x objective) were obtained with patch pipettes (2-5 Ω) pulled from borosilicate capillary glass with a Sutter P-97 horizontal puller. Pipettes contained, depend on experiment, one of the following solutions (in mM): *Cesium internal*: 140 Cs-gluconate, 5 HEPES, 1 MgATP, 10 phosphocreatine, 10 TEA-Cl, 1 EGTA. *Potassium internal*: 130 K-gluconate, 8 KCl, 1 MgCl₂, 3 Mg-ATP, 10 HEPES, 10 phosphocreatine, 1.1 or 0.2 EGTA. pH was adjusted to 7.2 with Cs-OH or K-OH. In a subset of re-patching experiments, different Alexa dyes (488 and 594) were included in the cesium and potassium filled pipettes to verify that the same pyramidal cell had been patched twice. Interneurons were patched in transgenic B13 and G42 mouse lines that selectively express GFP in parvalbumin expressing basket interneurons using epifluorescence.

Data acquisition and analysis

Data were recorded with a Multiclamp 700B amplifier, digitized at 10-20 kHz, and analyzed offline. In dual whole-cell recordings with a cesium internal solution, unclamped spikes ("action currents") were elicited in pyramidal cells under voltage clamp by applying brief (1 ms) depolarizing voltage pulse from the holding potential of -70 mV. With potassium internals, action potentials were generated by applying brief (5 ms) depolarizing current pulse. The presynaptic cells were stimulated at 0.1 Hz. IPSCs were recorded at the EPSC reversal potential

(0 mV), and were confirmed to be IPSCs by recording at the IPSC reversal potential (-70 mV). Average values are expressed as mean \pm SEM. The student's *t*-test was used for statistical comparisons, with significance defined as $p < 0.01$.

Latencies

The latency of postsynaptic currents was determined by the time difference between the peak of the presynaptic spike and the 5% rise point of postsynaptic currents. Despite the broad distribution of latencies between the spike evoked in a pyramidal cell and onset of IPSCs in a simultaneously recorded pyramidal cell (Fig. 1A), even the shortest latencies are consistent with a disynaptic IPSC: The average latency of the unitary EPSC from pyramidal cells to FS cells was 0.61 ± 0.04 ms ($n = 8$), and the minimal latency was 0.40 ms. The average latency of the unitary IPSC from FS cells to pyramidal cells was 0.62 ± 0.05 ms ($n = 8$), and the minimal latency was 0.40 ms. Given the fast membrane time constant and rapid kinetics of EPSPs at FS cells (<0.5 ms for 20-80% rise of EPSP (Geiger et al., 1997)), and that FS cells can integrate and fire within 0.6 ms (Gabernet et al., 2005), a latency as short as 1.8 ms is a reasonable time to expect for a disynaptic connection. Based on our minimal latencies of 0.4 ms for monosynaptic connections, however, even shorter latencies could be expected. Indeed, in our dataset, 6% were below 2 ms.

Estimating the Fractional Recruitment of Interneurons

Both Ren et al. and our data show average unitary IPSC (uIPSC) amplitudes of about 100 pA, and pyramidal cell evoked IPSC amplitudes of about 200 pA. Given a bi-directional connection probability between fast spiking interneurons and pyramidal cells of 0.5-0.75 (Holmgren et al., 2003), average IPSCs result from the recruitment of four or less interneurons. In order to estimate what fraction of connected interneurons this represents, we evaluated the density of interneurons within a conservative cylindrical cortical volume of 200 μ m in diameter (even though pyramidal cells are certain to contact even more interneurons outside of this volume). Based on a cortical neuron density of 100,000/mm³, such a volume would contain 628 pyramidal cells (Holmgren et al. 2003). While estimates of interneuron density vary (Beaulieu, 1993; Gonchar et al., 2007), if we assume 20% of layer 2/3 cells are interneurons, and 50% of these interneurons are basket cells (Markram et al., 2004), then we can estimate that there are 79 basket cells in this volume. Thus, a pyramidal cell would contact about 40-60 interneurons, and only 7-10% of these (4 out of 40 to 60) would need to be recruited to generate an IPSC.

In Utero Electroporation

Timed pregnant ICR white mice (Charles River) at E15-16 were operated on as previously described (Saito and Nakatsuji, 2001). Each embryo was injected with 1 μ g of pCAGGS-ChR2-Venus (Petreanu et al., 2007) mixed with 0.5 μ g of pCAG-EGFP.

ChR2 stimulation

ChR2-expressing layer 2/3 pyramidal cells were activated by a full-field 5 ms flashes of light from a 5 W luxeon blue LED coupled to the epifluorescence pathway of the Olympus BX51. To record synaptic currents in the presence of TTX, we bath applied the potassium channel blocker, 4-AP, at 1 mM (Petreanu et al. 2009).

Results

We performed whole-cell paired recordings from neighboring (<75 μ m) pyramidal cells in layer 2/3 of the rat somatosensory cortex. Consistent with previous findings (Kapfer et al., 2007, 0 of 305 connections; Silberberg and Markram, 2007, 4 of 1450 pairs), single action potentials triggered in one of the pyramidal cells failed to evoke disynaptic inhibitory

postsynaptic currents (IPSC) in the neighboring pyramidal cell (0 of 19). To test whether these results were specific to the somatosensory cortex, we repeated these experiments in the rat visual cortex. As in the somatosensory cortex, we could not evoke disynaptic IPSCs in one of the pyramidal cells in response to individual spikes triggered in the other (0 of 6). These results differ, however, from a recent study in the visual cortex of mice, which showed large IPSCs in layer 2/3 pyramidal cells evoked by triggering a single action potential in a neighboring pyramidal cell (Ren et al., 2007).

To test whether this difference is species specific between the cortex of rats and mice, we repeated these experiments in the mouse visual cortex. As in the rat, we could not evoke disynaptic IPSCs in layer 2/3 pyramidal cells in response to single spikes triggered in neighboring pyramidal cells of the mouse visual cortex (0 of 48). We did, however, observe monosynaptic excitatory postsynaptic currents (EPSCs) in 17% of recordings (8 of 48), consistent with the known connectivity between layer 2/3 pyramidal cells (connection probability of 0.1 to 0.2: Mason et al., 1991; Thomson and Deuchars, 1997; Atzori et al., 2001; Holmgren et al., 2003; Yoshimura et al., 2005; Feldmeyer et al., 2006; Kapfer et al., 2007). As in the visual cortex, we were also unable to evoke disynaptic IPSCs in the mouse somatosensory cortex using the same experimental protocol (0 of 6). Hence, in both the visual and somatosensory cortices of rats and mice, single action potentials evoked in layer 2/3 pyramidal cells did not evoke disynaptic IPSCs in neighboring pyramidal cells (0 of 79 total) as previously reported.

In the study published by Ren et al. (2007), disynaptic IPSCs were recruited by triggering action currents in pyramidal cells filled with a cesium-based internal solution (though a few IPSCs, 4 of 28 pairs, were generated using a potassium-based internal). Thus, to test whether this experimental difference could account for the discrepancy between our results, we performed paired recordings between pyramidal cells filled with a cesium based internal solution (fig. 1). Under these conditions, a brief depolarizing voltage step (40 mV for 1 ms) applied to a pyramidal cell held at -70 mV in voltage-clamp produced an un-clamped spike (or “action current”; fig. 1A). In 7 of 22 of these recordings (32%; 5 of 17 in mouse visual cortex and 2 of 5 in rat somatosensory cortex; the same percentage as observed by Ren et al.), the action current elicited large IPSCs in simultaneously recorded neighboring pyramidal cells (fig. 1A).

To ensure that the difference between the two series of experiments was indeed due to different internal solutions rather than to a sampling artifact, we performed a series of paired recordings where one of the two pyramidal cells was first patched with a potassium based internal solution, and subsequently re-patched with a cesium based solution (fig. 1B). While action potentials triggered in pyramidal cells dialyzed with a potassium based solution never evoked disynaptic IPSCs in neighboring pyramidal cells, re-patching the same pyramidal cell with a cesium based internal produced disynaptic IPSCs in 9 of 47 experiments (19%) in the mouse somatosensory cortex, rat visual cortex and rat somatosensory cortex (for all IPSCs recorded in cesium: mean amplitude: 190.2 ± 36.9 pA; mean latency: 7.1 ± 1.1 ms; $n = 16$; fig. 1B). Replacing potassium with cesium did not, however, produce EPSCs between pyramidal pairs that were not already monosynaptically coupled. These results suggests that a cesium based internal solution is required to facilitate the recruitment of inhibition in response to single spikes in layer 2/3 pyramidal cells.

To determine the mechanism that facilitates the recruitment of inhibition with a cesium based internal solution, we performed paired recordings between layer 2/3 pyramidal cells and fast-spiking (FS) inhibitory interneurons to compare the amplitude of monosynaptic unitary excitatory postsynaptic potentials (EPSPs) when the presynaptic pyramidal cell was dialyzed with a potassium or cesium solution (fig. 1C). Specifically, pyramidal cells were first patched

with a potassium based internal solution, and then re-patched with a cesium solution while maintaining the same postsynaptic FS interneuron. Switching from a potassium to a cesium based internal solution doubled the size of the unitary EPSPs recorded at FS interneurons (2.2 fold, $n=8$; fig. 1C). Furthermore, this doubling of EPSP size was directly attributable to the cesium internal, and not the voltage-clamp paradigm used to produce action currents. With a potassium internal solution, there was no difference in the amplitude of the unitary EPSP when spikes were generated in pyramidal cells with either current clamp or escape from voltage-clamp (*I-clamp*: 2.1 ± 0.6 mV, *V-clamp*: 2.1 ± 0.6 mV, $n = 6$; fig. 1D).

The increased unitary excitation of inhibitory interneurons provides a plausible mechanism for the observed disynaptic IPSPs in response to a single spike in a cesium dialyzed pyramidal cell, as it has been shown previously that doubling the unitary EPSP amplitude onto interneurons strongly increases their spiking probability (Kapfer et al., 2007). However, a cesium-mediated increase in synaptic transmission does not exclude the possibility suggested by Ren et al. (2007) that glutamate release may also directly activate presynaptic receptors located on GABAergic terminals and, as a consequence, trigger GABA release.

To directly address this possibility, we functionally isolated GABAergic axon terminals from the somatodendritic compartment by blocking action potentials with TTX. In order to trigger glutamate release from pyramidal cells in the presence of TTX, we transfected layer 2/3 pyramidal cells with channelrhodopsin-2 (ChR2) through *in utero* electroporation of pregnant mice. Importantly, this procedure does not transfect interneurons, and therefore selectively targets pyramidal cells. The expression of ChR2 along the axon of pyramidal cells allows glutamate release to be triggered through photo-stimulation even in the absence of action potentials (Petreanu et al. 2009).

Consistent with Petreanu et al. (2009), photo-stimulation of layer 2/3 pyramidal cell axons in somatosensory cortex in the presence of TTX and 4-AP (to prolong synaptic depolarization, Kole et al., 2007; Shu et al., 2007; Goldberg et al., 2008) could robustly evoke synaptic glutamate release, as shown by the large excitatory postsynaptic conductances recorded in pyramidal cells voltage clamped at -80 mV (8 ± 1 nS; $n=7$; fig. 2). Because the unitary EPSC conductance between two layer 2/3 pyramidal cells averages 0.24 nS (Kapfer et al., 2007), photostimulation triggered the release from approximately 30 axons converging on the recorded pyramidal cell. To determine whether glutamate released from these photostimulated axons triggered the disynaptic release of GABA from inhibitory terminals, we voltage-clamped the recorded pyramidal cell at the reversal potential of EPSCs, around 0 mV to isolate IPSCs. Photostimulation never evoked an IPSC (0 of 20) as illustrated by the complete absence of an outward current in pyramidal cells (fig. 2). We did not detect an IPSC under these conditions for stimulation of layer 2/3 of somatosensory cortex ($n=7$; fig. 2) or visual cortex ($n=13$).

Hence, under the present conditions glutamate release from pyramidal cells does not trigger GABA release by directly acting on GABAergic terminals. Rather, our data support the idea that for layer 2/3 inhibitory circuitry GABA release occurs in response to action potentials generated by classical somatodendritic integration of EPSPs.

Discussion

In agreement with the report of Ren et al. (2007), the present data show that when layer 2/3 pyramidal cells of primary sensory cortices are recorded with a cesium containing intracellular solution, a single action current in a pyramidal cell is sufficient to trigger IPSCs at a latency consistent with disynaptic recruitment of inhibition (see Methods) in approximately 30% of neighboring pyramidal cells. We show that this apparent contradiction with previous observations of ours and other laboratories (i.e. where a single spike in a pyramidal cell never

or in extremely rare cases triggers inhibition in its neighbors, e.g. Mason et al., 1991; Deuchars et al., 1994; Kapfer et al., 2007; Silberberg and Markram, 2007; Lefort et al., 2009) is likely to be due to the presence of cesium rather than potassium in the presynaptic recording pipette. Cesium, a potassium channel impermeant ion, can increase the probability of synaptic vesicle release by dramatically prolonging the depolarization at a presynaptic terminal during an action potential (Kole et al., 2007). Indeed, we show that intracellular cesium more than doubles the size of all unitary EPSPs, regardless of size, evoked by a pyramidal cell onto inhibitory interneurons.

Because of the high connectivity between interneurons and pyramidal cells (Holmgren et al., 2003; Kapfer et al., 2007), even the spiking of very few (1-2) interneurons in the entire slice has a high probability of being detected through an IPSC in a randomly patched pyramidal cell. The doubling of EPSP amplitudes should recruit these few interneurons in a conventional layer 2/3 disynaptic circuit. Though our data provide only a small sample of the distribution of EPSP sizes from pyramidal cells onto interneurons, previous work has shown that a significant fraction (>10%) of these inputs are between 5 and 12 mV (Holmgren et al., 2003). Doubling the size of these large EPSPs to 10-24 mV would almost certainly bring some fraction of interneurons above threshold. The pyramidal cell evoked IPSCs we describe, as well as those from Ren et al., are about twice the amplitude of uIPSCs. Given a connection probability from FS cells to pyramidal cells of 0.5-0.75, this suggests that IPSCs result from the recruitment of at most only four inhibitory interneurons, which corresponds to only about 7-10% of the basket cells contacted by any single pyramidal cell a cylindrical volume of cortex 200 μ m in diameter (Beaulieu, 1993; Holmgren et al., 2003; Markram et al., 2004; Gonchar et al., 2007; see Methods). Unfortunately, this sparse activation of interneurons also means that direct identification of the few spiking cells will require finding the proverbial needle in a very large haystack of cortical neurons.

Contrary to the expectation that large EPSPs will spike some interneurons, Ren et al. (2007) make the provocative suggestion that at least part of the observed disynaptic inhibition of pyramidal cells is due to the direct action of released glutamate onto GABAergic terminals, rather than on the somatodendritic compartment of inhibitory interneurons. Furthermore, this mode of evoked GABA release does not appear involve the recruitment of action potentials in the inhibitory axon. If true, this would allow for individual pyramidal cells to trigger disynaptic inhibition in a highly localized manner, defined by the arborization of their own axon rather than by the arborization of the recruited interneurons. To test this possibility, we used TTX to functionally uncouple the axon and somatodendritic compartment of GABAergic interneurons while evoking glutamate release via photostimulation of pyramidal cells expressing channelrhodopsin-2. Photostimulation triggered very large excitatory currents, due to the release of glutamate from many axons. However, photo-released glutamate failed to trigger GABA release, as shown by the complete absence of inhibitory currents in the recorded pyramidal cells.

Our data are thus inconsistent with a circuit involving GABA release via direct activation of glutamate receptors on presynaptic terminals of GABAergic interneurons by synaptically released glutamate. We do not, however, exclude the possibility that exogenous application of glutamatergic agonists may lead to the activation of GABAergic terminals, as shown by Ren et al. (2007). While we believe that the presence of presynaptic cesium can explain most of the phenomena reported by Ren et al., unfortunately there are a number of inconsistencies that we are unable to replicate, including the recruitment of IPSCs with a potassium internal, the rare IPSCs with near-millisecond latencies, and the recruitment of inhibition in TTX. Our data support the hypothesis that, at least in layer 2/3 of the cortex, disynaptic inhibition results from the classical integrate and fire behavior of inhibitory interneurons.

Acknowledgements

We are grateful to L. Petreanu and K. Svoboda for suggesting the photostimulation experiment. We thank J. Isaacson, L. Glickfeld, B. Atallah, F. Pouille and A. Marin-Burgin for their comments on the manuscript. This work was supported by NIH Grant #NH70058. CH was supported by a post-doctoral NRSA fellowship (5F32NS060585-03).

References

- Atzori M, Lei S, Evans DI, Kanold PO, Phillips-Tansey E, McIntyre O, McBain CJ. Differential synaptic processing separates stationary from transient inputs to the auditory cortex. *Nat Neurosci* 2001;4:1230–1237. [PubMed: 11694887]
- Beaulieu C. Numerical data on neocortical neurons in adult rat, with special reference to the GABA population. *Brain Res* 1993;609:284–292. [PubMed: 8508310]
- Chattopadhyaya B, Di Cristo G, Higashiyama H, Knott GW, Kuhlman SJ, Welker E, Huang ZJ. Experience and activity-dependent maturation of perisomatic GABAergic innervation in primary visual cortex during a postnatal critical period. *J Neurosci* 2004;24:9598–9611. [PubMed: 15509747]
- Connors BW, Cruikshank SJ. Bypassing interneurons: inhibition in neocortex. *Nat Neurosci* 2007;10:808–810. [PubMed: 17593940]
- Deuchars J, West DC, Thomson AM. Relationships between morphology and physiology of pyramidal single axon connections in rat neocortex in vitro. *J Physiol* 1994;478(Pt 3):423–435. [PubMed: 7965856]
- Dumitriu D, Cossart R, Huang J, Yuste R. Correlation between axonal morphologies and synaptic input kinetics of interneurons from mouse visual cortex. *Cereb Cortex* 2007;17:81–91. [PubMed: 16467567]
- Feldmeyer D, Lubke J, Sakmann B. Efficacy and connectivity of intracolumnar pairs of layer 2/3 pyramidal cells in the barrel cortex of juvenile rats. *J Physiol* 2006;575:583–602. [PubMed: 16793907]
- Gabernet L, Jadhav SP, Feldman DE, Carandini M, Scanziani M. Somatosensory integration controlled by dynamic thalamocortical feed-forward inhibition. *Neuron* 2005;48:315–327. [PubMed: 16242411]
- Geiger JR, Lubke J, Roth A, Frotscher M, Jonas P. Submillisecond AMPA receptor-mediated signaling at a principal neuron-interneuron synapse. *Neuron* 1997;18:1009–1023. [PubMed: 9208867]
- Goldberg EM, Clark BD, Zagha E, Nahmani M, Erisir A, Rudy B. K⁺ channels at the axon initial segment dampen near-threshold excitability of neocortical fast-spiking GABAergic interneurons. *Neuron* 2008;58:387–400. [PubMed: 18466749]
- Gonchar Y, Wang Q, Burkhalter A. Multiple distinct subtypes of GABAergic neurons in mouse visual cortex identified by triple immunostaining. *Front Neuroanat* 2007;1:3. [PubMed: 18958197]
- Holmgren C, Harkany T, Svennenfors B, Zilberter Y. Pyramidal cell communication within local networks in layer 2/3 of rat neocortex. *J Physiol* 2003;551:139–153. [PubMed: 12813147]
- Howard A, Tamas G, Soltesz I. Lighting the chandelier: new vistas for axo-axonic cells. *Trends Neurosci* 2005;28:310–316. [PubMed: 15927687]
- Kapfer C, Glickfeld LL, Atallah BV, Scanziani M. Supralinear increase of recurrent inhibition during sparse activity in the somatosensory cortex. *Nat Neurosci* 2007;10:743–753. [PubMed: 17515899]
- Kole MHP, Letzkus JJ, Stuart GJ. Axon Initial Segment Kv1 Channels Control Axonal Action Potential Waveform and Synaptic Efficacy. *Neuron* 2007;55:633–647. [PubMed: 17698015]
- Lefort S, Tómm C, Floyd Sarria JC, Petersen CC. The excitatory neuronal network of the C2 barrel column in mouse primary somatosensory cortex. *Neuron* 2009;61:301–316. [PubMed: 19186171]
- Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C. Interneurons of the neocortical inhibitory system. *Nat Rev Neurosci* 2004;5:793–807. [PubMed: 15378039]
- Mason A, Nicoll A, Stratford K. Synaptic transmission between individual pyramidal neurons of the rat visual cortex in vitro. *J Neurosci* 1991;11:72–84. [PubMed: 1846012]
- Petreanu L, Huber D, Sobczyk A, Svoboda K. Channelrhodopsin-2-assisted circuit mapping of long-range callosal projections. *Nat Neurosci* 2007;10:663–668. [PubMed: 17435752]
- Petreanu L, Mao T, Sternson S, Svoboda K. The subcellular organization of neocortical excitatory connections. *Nature*. 2009(in press)
- Ren M, Yoshimura Y, Takada N, Horibe S, Komatsu Y. Specialized inhibitory synaptic actions between nearby neocortical pyramidal neurons. *Science* 2007;316:758–761. [PubMed: 17478724]

- Rudomin P. Presynaptic inhibition of muscle spindle and tendon organ afferents in the mammalian spinal cord. *Trends Neurosci* 1990;13:499–505. [PubMed: 1703681]
- Saito T, Nakatsuji N. Efficient gene transfer into the embryonic mouse brain using in vivo electroporation. *Dev Biol* 2001;240:237–246. [PubMed: 11784059]
- Schoppa NE, Urban NN. Dendritic processing within olfactory bulb circuits. *Trends Neurosci* 2003;26:501–506. [PubMed: 12948662]
- Sherman SM. Interneurons and triadic circuitry of the thalamus. *Trends Neurosci* 2004;27:670–675. [PubMed: 15474167]
- Shu Y, Yu Y, Yang J, McCormick DA. Selective control of cortical axonal spikes by a slowly inactivating K⁺ current. *Proc Natl Acad Sci U S A* 2007;104:11453–11458. [PubMed: 17581873]
- Silberberg G, Markram H. Disynaptic inhibition between neocortical pyramidal cells mediated by Martinotti cells. *Neuron* 2007;53:735–746. [PubMed: 17329212]
- Thomson AM, Deuchars J. Synaptic interactions in neocortical local circuits: dual intracellular recordings in vitro. *Cereb Cortex* 1997;7:510–522. [PubMed: 9276176]
- Wassle H. Parallel processing in the mammalian retina. *Nat Rev Neurosci* 2004;5:747–757. [PubMed: 15378035]
- Yoshimura Y, Dantzker JL, Callaway EM. Excitatory cortical neurons form fine-scale functional networks. *Nature* 2005;433:868–873. [PubMed: 15729343]




Figure 1. Increasing glutamate release in individual pyramidal cells with cesium dialysis promotes disynaptic inhibition

A. IPSCs can be generated between pyramidal cell pairs in L2/3 with a cesium based internal solution. *Left:* A paired recording from two nearby L2/3 pyramidal cells with a cesium internal solution in both cells. A brief depolarizing voltage step (40 mV for 1 ms) applied to one pyramidal cell (top) yields an un-clamped spike (or “action current”). This spike produces a large IPSC in a neighboring pyramidal cell (bottom, 0 mV). *Right:* Latencies and amplitudes of IPSCs generated with cesium internal solutions. IPSCs were observed in 16 of 69 (23%) total recordings with a cesium internal (7 of 22, or 32% when both cells are initially filled with cesium, 9 of 47, or 19% when one cell is patched twice (fig 2.) and 0 of 79 (0%) recordings

with a potassium internal solution. IPSCs were generated with a cesium internal in the mouse visual cortex, mouse somatosensory (ss) cortex, rat visual cortex and rat somatosensory (ss) cortex.

B. Repatching with a cesium based internal solution generates IPSCs between pyramidal cell pairs. *Left*: A pair of pyramidal cells, in which one cell contains a potassium internal (top) and the other contains a cesium internal (bottom; 0 mV). Action potentials from the pyramidal cell in potassium do not produce IPSCs in the pyramidal cell filled with cesium. *Right*: when the same pyramidal cell is re-patched with a cesium based internal solution, spikes produced by depolarizing voltage steps (as in panel A) now produce IPSCs in the neighboring paired pyramidal cell. Repatching in this manner produced IPSCs in 9 of 47 (19%) recordings from the mouse somatosensory (ss) cortex, rat visual cortex and rat somatosensory (ss) cortex.

C. Repatching pyramidal cells with a cesium based internal solution doubles the size of EPSPs onto fast-spiking (FS) interneurons. Paired recordings were performed between layer 2/3 pyramidal cells and fast-spiking (FS) interneurons. In these experiments, pyramidal cells were first patched with a potassium internal solution in order to measure the amplitude of unitary EPSPs onto FS interneurons. Pyramidal cells were then re-patched with a cesium internal solution to re-measure the unitary EPSPs in response to spikes produced by depolarizing voltage steps (as in panel A). Switching from a potassium to a cesium based internal solution produced a 2.2 fold increase in FS cell EPSPs on average (n=8).

D. The increase in FS interneuron EPSPs does not result from the voltage-clamp paradigm *per se*. With a potassium internal solution, there is no difference in the amplitude of FS interneuron EPSPs when spikes are generated in pyramidal cells with either current clamp or escape from voltage-clamp.




Figure 2. Strong glutamate release from pyramidal cells does not drive action-potential independent disynaptic inhibition

A. Brief flashes of blue light to ChR2-expressing L2/3 pyramidal cells drives neurotransmitter release onto neighboring ChR2-negative L2/3 cells. Cells exhibit both glutamatergic ($V_h = -70$ mV) and disynaptic GABAergic ($V_h = 0$ mV) responses (note temporal delay in inset) in the absence of TTX (left), but only direct glutamatergic responses when action potentials are blocked and synaptic release is facilitated by the potassium channel blocker 4-AP.

B. Example experiment showing the amplitude of the post-synaptic current before and during TTX application, and in the additional presence of 4-AP. The cell is held at -70 mV except as indicated.

C. Left: Amplitude of post-synaptic currents recorded at -70 and 0 mV in the presence of TTX and 4-AP in layer 2/3 of somatosensory cortex (mean at -70 mV = -653 ± 75 pA, at 0 mV = -5 ± 5 pA, n = 7). Right: Currents for cells recorded in visual cortex (mean at -70 mV = -780 ± 156 pA, at 0 mV = 4 ± 3 pA, n = 13).